

Gill SM

National Aeronautics and Space Administration
Goddard Space Flight Center
Contract No. NAS-5-9299

ST - PF - 10490

EFFECT OF RADIATION ATTENUATION
UPON THE MOTION OF A RELATIVISTIC PARTICLE
IN A UNIFORM FIELD

by

G. E. Gernet

(USSR)

FACILITY FORM 602

| | |
|-------------------------------|------------|
| N 67 12203 | (THRU) |
| (ACCESSION NUMBER) | 1 |
| 5 | (CODE) |
| (PAGES) | 24 |
| NR 80179 | (CATEGORY) |
| (NASA CR OR TMX OR AD NUMBER) | |

GPO PRICE \$ _____

CFSTI PRICE(S) \$ _____

Hard copy (HC) 1.00

Microfiche (MF) 50

ff 653 July 65

20 MAY 1966

EFFECT OF RADIATION ATTENUATION
UPON THE MOTION OF A RELATIVISTIC PARTICLE
IN A UNIFORM MAGNETIC FIELD*

Doklady AN SSSR, Fizika,
 Tom 168, No. 1, 63-4,
 Izdatel'stvo "NAUKA", 1966

by G. E. Gernet

SUMMARY

The effect of radiation attenuation upon the motion of a relativistic particles in a uniform magnetic field is evaluated by the behavior of the transverse velocity components, for which the approximate equations are derived.

* * *

When a charged particle moves in a magnetic field, its energy decreases because of magnetic bremsstrahlung (synchrotron) radiation, and this leads to trajectory change.

In the cases, when the particle's sojourn in the magnetic field is comparable with the characteristic time of energy decrease

$$T \sim m^3 c^5 / e^4 H^2$$

(as, for example, for electrons in cosmic fields), one must take into account the radiation attenuation when determining the trajectory.

The particle's equation of motion has the form

$$\frac{dp}{dt} = \frac{e}{c} [\mathbf{v}, \mathbf{H}] + \mathbf{f}_r, \quad (1)$$

where the attenuation force \mathbf{f}_r is [1]

$$\mathbf{f}_r = \frac{2e^4}{3m^2 c^5} \left\{ [\mathbf{H} [\mathbf{H}, \mathbf{v}]] - \frac{\mathbf{v}}{1 - v^2/c^2} \frac{1}{c^2} ([\mathbf{v}, \mathbf{H}])^2 \right\}. \quad (2)$$

Let us direct the axis \underline{z} of the Descartes system of coordinates

* VLIYANIYE RADIATIONNOGO TORMOZHENIYA NA DVIZHENIYE RELYATIVISTKOY CHASTITSY V ODNORODNOM MAGNITNOM POLE

along the field and denote

$$\omega = eH / mc, \quad \delta = 2/3 e^4 H^2 / m^3 c^5. \quad (3)$$

We shall express v in fractions of c and the energy E in fractions of mc^2 :

$$u = v / c, \quad w = E / mc^2 = 1 / \sqrt{1 - u^2}.$$

Passing then to components and taking into account that $p = Ev / c^2$, we shall obtain

$$\begin{aligned} du_x w / dt &= \omega u_y - \delta u_x (1 - u_z^2) w^2, \\ du_y w / dt &= -\omega u_x - \delta u_y (1 - u_z^2) w^2, \\ du_z w / dt &= -\delta u_z w^2 (u_x^2 + u_y^2). \end{aligned} \quad (4)$$

It follows from (4), first of all, that

$$u_z = \text{const.} \quad (5)$$

Taking into account (5) we obtain from (4) the equation for energy variation

$$dw / dt = -\delta [(w / w_\infty)^2 - 1], \quad (6)$$

where

$$w_\infty = 1 / \sqrt{1 - u_z^2} \quad (7)$$

is the limit value of w at $t \rightarrow \infty$.

The integral of (6) is

$$w = w_\infty \text{cth} (\delta t / w_\infty + C_0). \quad (8)$$

The constant C_0 is determined from the condition $w = w_0$ at $t = 0$.

It is appropriate to express the quantity w_∞ by the angle θ between the initial velocity and the direction of the field and by the initial value of energy. Assuming in (7) $u_z = u_0 \cos \theta$, and taking into account that $w_0 = 1 / \sqrt{1 - u_0^2}$, we have

$$w_\infty = \frac{w_0}{\sqrt{\cos^2 \theta + u_0^2 \sin^2 \theta}}. \quad (9)$$

Taking advantage of (8), we may integrate the equations for the transverse components u_x and u_y

$$\begin{aligned} u_x &= u_\perp(0) e^{-\delta \tau} \sin (\omega \tau + \varphi_0), \\ u_y &= u_\perp(0) e^{-\delta \tau} \cos (\omega \tau + \varphi_0), \end{aligned} \quad (10)$$

where

$$u_{\perp}(0) = \sqrt{u_x^2(0) + u_y^2(0)};$$

φ_0 is the initial phase; τ is the proper time

$$\tau = \int_0^t \frac{dt}{w} = \frac{1}{\delta} \ln \frac{\text{ch}(\delta t/w_{\infty} + C_0)}{\text{ch} C_0}. \quad (11)$$

It may be seen from (10) that the transverse velocity components damp and become zero at $t \rightarrow \infty$.

The proper time is a complex function of t , as distinct from the motion without taking into account the radiation friction, when $\tau = t/w_0$. As a consequence of that the variation of u_x and u_y with time will no longer be harmonic, which should exert an influence on the emission spectrum.

The equations obtained are substantially simplified when passing to the extreme relativistic case characterized by the correlation $w_0 \gg 1$.

Outside a narrow cone around the direction with an angle $\theta_0 \sim 1/w_0^2$ we have from (9) $w_{\infty} \approx 1/\sin \theta$. Then (8) passes into

$$\frac{1}{w} = \frac{1}{w_0} + \sin \theta \text{th}(\delta t \sin \theta). \quad (12)$$

Inasmuch as $\text{th } x$ practically attains its limit value at a value of x slightly greater than the unity, the main energy loss will take place in a time

$$t_0 \sim 1/\delta \sin \theta. \quad (13)$$

In the interval $0 \leq t \leq t_0$ it is sufficient to limit oneself to the first term of the expansion of $\text{th } x$ in series, so that

$$\frac{1}{w} = \frac{1}{w_0} + (\sin^2 \theta) \delta t \quad (t \delta \sin \theta \ll 1). \quad (14)$$

Hence it may be seen that the energy decreases by a factor of two in a time $\sim 1/w_0 \delta \sin^2 \theta$.

For the proper time we have in the same approximation

$$\tau = \frac{t}{w_0} + t^2 \frac{\delta}{2} \sin^2 \theta. \quad (15)$$

Substituting (15) into (10), we shall obtain the approximate expressions for the transverse velocity components.

I am grateful to Prof. L. E. Gurevich for discussing the work.

*** THE END ***

Leningrad Electr. Engin. Institute
of Communications in the name of
M. A. Bonch-Bruyevich

Received on
3 June 1965.

REFERENCE

[1].- L. D. LANDAU, E. M. LIFSHITS.- Teoreticheskaya fizika, 2, ch. IX, 76, M
1960.

Consultants and Designers, Inc.
Arlington, Virginia
Contract No. NAS-5-9299

Translated by ANDRE L. BRICHANT
on 20 May 1966

DISTRIBUTION

GOODARD SPACE F.C.

NASA HQS

OTHER CENTERS

100 CLARK, TOWNSEND
110 STROUD
400 BOURDEAU
610 MEREDITH
611 McDONALD
ABRAHAM BRUNSTEIN
DAVIS EVANS
WILLIAMS
KONRADI
BOLDT VKB
SERLEMITOS
612 HEPPNER
NESS
SKILLMAN
613 KUPPERIAN
HALLAM
614 WHITE
KASTNER
615 BAUER
GOLDBERG HERMAN
MAIER STONE
640 HESS
MEAD MAEDA
SPEISER
NORTHROP
STERN
630 GI for SS [3]
252 FIBRARY
256 FREAS

SS NEWELL, NAUGLE
SG MITCHELL
SCHARDT
SCHMERLING
DUBIN
SL LIDDEL
FELLOWS
HIPSHER
HOROWITZ
SM FOSTER
GILL
RR KURZWEG
RTR NEILL
ATSS - T
WX SWEET

AMES R C
SONETT
LIBRARY
LANGLEY R C
160 ADAMSON
185 WEATHERWAX
JPL
SNYDER
NEUGEBAUER
WYCKOFF
U. IOWA
VAN ALLEN
UCLA
COLEMAN
UC BERKELEY
WILCOX
SWCAS
JOHNSON